

LEVY SOLUTIONS OF A RANDOMLY FORCED BURGERS EQUATION

MARIE-LINE CHABANOL AND JEAN DUCHON

ABSTRACT. We consider Burgers equation forced by a brownian in space and white noise in time process $\partial_t u + \frac{1}{2}\partial_x(u)^2 = f(x, t)$, with $E(f(x, t)f(y, s)) = \frac{1}{2}(|x| + |y| - |x - y|)\delta(t - s)$ and we show that there are Levy processes solutions, for which we give the evolution equation of the characteristic exponent. In particular we give the explicit solution in the case $u_0(x) = 0$.

1. INTRODUCTION

We consider the randomly forced Burgers equation $\partial_t u + \frac{1}{2}\partial_x(u)^2 = f(x, t)$, where f satisfies $E(f(x, t)f(y, s)) = A(x, y)\delta(t - s)$ with $A(x, y) = \frac{1}{2}(|x| + |y| - |x - y|)$: f is a brownian in space/white noise in time process (the brownian is considered on the whole space, and is obtained with two independent brownians, one on the left and one on the right). Burgers equation has originally been introduced as a 1D model of turbulence. It is certainly too crude, but it is still a good idea when trying to find new approaches to the “real” 3D turbulence as given by Euler equation, to start by looking at Burgers. In this point of view, the randomly forced Burgers equation is a crude simplification on the way to the description of forced turbulence. A variant of our problem was considered by [8], where they considered a gaussian in space/white noise in time forcing, satisfying $E(f(x, t)f(y, s)) = A(x - y)\delta(t - s)$ and obtained all the hierarchy of n -point densities evolution equations. We will proceed quite differently, basing our study on the result shown by Carraro and Duchon in [6], and by Bertoin in [3], that Levy processes with negative jumps are statistical solutions (in a sense that we will recall later) of the Burgers equation without any forcing. This is actually a particular case of the fact that the Burgers equation conserves the Markovian (in space) property of processes ([7]). We will show here in the same spirit that the brownian in space/white noise in time forcing allows to keep the Levy property. Moreover, our method allows us to write explicitly the exponent of the Levy process that one gets when starting from $u_0(x) = 0$.

2. STATISTICAL SOLUTIONS

We will closely follow [6] (see also [13]). Let E be the space of càdlàg real functions. We will call $\mathcal{C}(E)$ the smallest σ -algebra such that for each $x \in \mathbb{R}$, $u \mapsto u(x)$ is measurable, and $\mathcal{C}'(E)$ the smallest σ -algebra such that for each $(x, y) \in \mathbb{R} \times \mathbb{R}$, $u \mapsto u(x) - u(y)$ is measurable. Let $\mathcal{D}_0$ be the set of real C^∞ functions v with compact support such that $\int_{\mathbb{R}} v(x)dx = 0$. A probability μ on $(E, \mathcal{C}(E))$ is then characterized by its characteristic function

$$v \in \mathcal{D} \mapsto \hat{\mu}(v) := \int_E \exp \left[i \int_{\mathbb{R}} u(x)v(x)dx \right] d\mu(u)$$

whereas a probability μ on $(E, \mathcal{C}'(E))$ is characterized by its characteristic function $v \in \mathcal{D}_0 \mapsto \hat{\mu}(v)$.

Let $u_0 : (\Omega, \mathcal{A}, P) \rightarrow E$ be a random process, defined on some probability space, and let $\mu_0 : \mathcal{C}(E) \rightarrow [0, 1]$ denote its probability law : u_0 will be our initial condition. We will let u evolve according to the non forced Burgers equation, obtaining thus a family of processes u_t where $u_t(x) = u(x, t)$ is the solution of Burgers at time t . Then it is not difficult to check that the

characteristic function $\hat{\mu}_t$ of u_t verifies :

$$\begin{aligned}
 \partial_t \hat{\mu}_t(v) &= \partial_t \left(\int_E \exp \left[i \int_{\mathbb{R}} u(x) v(x) dx \right] d\mu_t(u) \right) \\
 &= \int_E \partial_t \left\{ \exp \left[i \int_{\mathbb{R}} u(x, t) v(x) dx \right] \right\} d\mu_0(u_0) \\
 &= \int_E \exp \left[i \int_{\mathbb{R}} u(x, t) v(x) dx \right] \left(\partial_t \left[i \int_{\mathbb{R}} u(x, t) v(x) dx \right] \right) d\mu_0(u_0) \\
 (1) \quad &= \int_E \exp \left[i \int_{\mathbb{R}} u(x) v(x) dx \right] \left[i \int_{\mathbb{R}} \frac{1}{2} u(x)^2 v'(x) dx \right] d\mu_t(u)
 \end{aligned}$$

This motivated the definition given in [6] of a statistical solution of Burgers equation, as a family of random processes u_t whose characteristic functions are solutions of (1).

Here we will take for u_0 a Levy process with negative jumps, that is a càdlàg process with stationary homogenous increments and negative jumps, and we are really interested only in the law of the increments. Hence u_0 really defines a probability law on $(E, \mathcal{C}'(E))$. But $u \mapsto \int_{\mathbb{R}} u(x)^2 v'(x) dx$ is not $\mathcal{C}'(E)$ measurable, hence this evolving equation for $\hat{\mu}$ makes no sense. One way around this is to take a different approach and to work with the Hopf-Cole construction as in [3]. Another way was explored in [6] : it is based on the notion of *intrinsic statistical solution*, and involves working with $u(x, t) - \frac{1}{b-a} \int_a^b u(x, t) dx$ when $b - a$ gets large :

Definition 1. *An intrinsic statistical solution of Burgers equation is a set $(\mu_t)_{t \geq 0}$ of probabilities on $(E, \mathcal{C}'(E))$ such that for any $v \in \mathcal{D}_0$,*

$$\partial_t \hat{\mu}_t(v) = \frac{i}{2} \lim_{a \rightarrow -\infty, b \rightarrow +\infty} \int_E \left(\int_{\mathbb{R}} \left(u(x) - \frac{1}{b-a} \int_a^b u(y) dy \right)^2 v'(x) dx \right) \exp \left[i \int_{\mathbb{R}} u v \right] d\mu_t(u)$$

In order to work with Levy processes we will need their characterization by means of their exponent.

Definition 2. *If u is a Levy process, its characteristic exponent ψ is defined by $\forall x < y, \forall \lambda \in \mathbb{R}$,*

$$(2) \quad E\{\exp i[\lambda(u(y) - u(x))]\} = \exp[(y - x)\psi(\lambda)]$$

If the process is of finite variance with negative jumps, one can use analytic continuation to define its Laplace exponent $\phi(\lambda) := \psi(-i\lambda)$ as a function on $\mathbb{R}^+$.

In [6] it is shown that if u_0 is a Levy process of finite variance such that $\phi'_0(0) \geq 0$, and if we let u evolve according to the non forced Burgers equation, u_t is still a Levy process; its characteristic exponent ψ_t satisfies the evolution equation $\partial_t \psi = i \partial_\lambda(\psi^2)$.

Let now f be the random forcing, independent on u_0 . We will denote by μ_F its probability law. Restricted to events involving a fixed time, μ_F is the law of a brownian motion defined on $\mathbb{R}$ by gluing one brownian on $\mathbb{R}^+$ and the reflection of another independent brownian on $\mathbb{R}^-$. Such a brownian on $\mathbb{R}$ is a continuous process with independent homogenous increments null at 0.

The basic idea now is the following : if u_t is a Levy process with probability on $\mathcal{C}'(E)$ μ_t , and if we let u_t evolve according to our forced Burgers equation, $u(x, t + dt)$ will be the sum of $u_{NF}(x, t + dt)$ and $\sqrt{dt}B(x)$, where u_{NF} is the process we would have obtained without any forcing, and $B(x)$ is a brownian motion independent of u_{NF} . $u(x, t + dt)$ should thus still be a process with homogenous independent increments, that is, a Levy process with exponent the sum of the exponents. Since the characteristic exponent of a brownian is $\frac{x^2}{2}$, and since the evolution equation for the characteristic exponent in case of no forcing is $\partial_t \psi(t, w) = i \partial_w \psi(t, w) \psi(t, w)$, the evolution equation of the characteristic exponent of μ with random forcing should be $\partial_t \psi(t, w) = i \partial_w(\psi^2) + \frac{w^2}{2}$.

One can also get this result in a more rigorous way by working with characteristic functions. Let us first define statistical solutions for the forced Burgers equation. Let $u(x, t)$ be a (weak) solution of our forced Burgers equation with $u(., 0) = u_0$, $u(., t) \in E$ for $t > 0$. Let μ_t denote the

law of $u(\cdot, t)$ on $\mathcal{C}(E)$. Using Ito's formula, one then gets for each v in $\mathcal{D}$:

$$\begin{aligned}
 \partial_t \hat{\mu}_t(v) &= \int_E \partial_t \left\{ \exp \left[i \int_{\mathbb{R}} u(x) v(x) dx \right] \right\} d\mu_t(u) \\
 &= \int_F \int_E \partial_t \left\{ \exp \left[i \int_{\mathbb{R}} u(x, t) v(x) dx \right] \right\} d\mu_0(u_0) d\mu_F \\
 &= \int_F \int_E \exp \left[i \int_{\mathbb{R}} u(x, t) v(x) dx \right] \left(\partial_t \left[i \int_{\mathbb{R}} u(x, t) v(x) dx \right] \right. \\
 &\quad \left. - \frac{1}{2} \int_{\mathbb{R} \times \mathbb{R}} v(x) v(y) \langle f(x, t) f(y, t) \rangle_t dx dy \right) d\mu_0(u_0) d\mu_F \\
 &= \int_E \left(\int_{\mathbb{R}} i \frac{1}{2} u(x)^2 v'(x) dx - \frac{1}{2} \int_{\mathbb{R} \times \mathbb{R}} v(x) v(y) A(x, y) dx dy \right) \exp \left[i \int_{\mathbb{R}} u v \right] d\mu_t(u)
 \end{aligned}$$

Hence our definitions :

Definition 3. A statistical solution of the forced Burgers equation (*) is a set $(\mu_t)_{t \geq 0}$ of probabilities on $(E, \mathcal{C}(E))$ such that for any $v \in \mathcal{D}$,

$$\partial_t \hat{\mu}_t(v) = \int_E \left(\int_{\mathbb{R}} i \frac{1}{2} u(x)^2 v'(x) dx \right) \exp \left[i \int_{\mathbb{R}} u v \right] d\mu_t(u) - \left(\frac{1}{2} \int_{\mathbb{R} \times \mathbb{R}} v(x) v(y) A(x, y) dx dy \right) \hat{\mu}_t(v)$$

Definition 4. An intrinsic statistical solution of the forced Burgers equation (*) is a set $(\mu_t)_{t \geq 0}$ of probabilities on $(E, \mathcal{C}'(E))$ such that for any $v \in \mathcal{D}_0$,

$$\begin{aligned}
 \partial_t \hat{\mu}_t(v) &= \lim_{a \rightarrow -\infty, b \rightarrow +\infty} \int_E \left(\int_{\mathbb{R}} \frac{i}{2} \left(u(x) - \frac{1}{b-a} \int_a^b u(y) dy \right)^2 v'(x) dx \right) \exp \left[i \int_{\mathbb{R}} u v \right] d\mu_t(u) \\
 (3) \quad &- \left(\frac{1}{2} \int_{\mathbb{R} \times \mathbb{R}} v(x) v(y) A(x, y) dx dy \right) \hat{\mu}_t(v)
 \end{aligned}$$

One can remark that for $A(x, y) = \frac{1}{2}(|x| + |y| - |x - y|)$ and $v \in \mathcal{D}$, $\int_{\mathbb{R} \times \mathbb{R}} v(x) v(y) A(x, y) dx dy = \int_0^{+\infty} (w(x)^2 + (w(-\infty) - w(-x))^2) dx$ where $w(x) = \int_x^{+\infty} v(y) dy$. If moreover $v \in \mathcal{D}_0$, $\int_{\mathbb{R} \times \mathbb{R}} v(x) v(y) A(x, y) dx dy = \int_{\mathbb{R}} w(x)^2 dx$.

A justification for this definition is the proposition, proven in [6], that if u is a homogeneous process of finite variance, then

$$\begin{aligned}
 &\lim_{a \rightarrow -\infty, b \rightarrow +\infty} \int_E \left(\int_{\mathbb{R}} \frac{i}{2} \left(u(x) - \frac{1}{b-a} \int_a^b u(y) dy \right)^2 v'(x) dx \right) \exp \left[i \int_{\mathbb{R}} u v \right] d\mu(u) = \\
 &\int_E \left(\int_{\mathbb{R}} i \frac{1}{2} u(x)^2 v'(x) dx \right) \exp \left[i \int_{\mathbb{R}} u v \right] d\mu(u)
 \end{aligned}$$

3. LEVY SOLUTIONS

3.1. Evolution equation. Now we can look for solutions that would be Levy processes with finite variance. Such processes are characterized by their exponent ψ_t , or by their Laplace exponent ϕ_t .

Let us recall here a useful lemma characterizing Laplace exponents of Levy processes.

Lemma 1. A function $\phi : \mathbb{R}^+ \mapsto \mathbb{R}$ is the Laplace exponent of a homogeneous Levy process with finite variance and negative jumps if and only if ϕ is C^∞ on $]0, +\infty[$, C^2 on $\mathbb{R}^+$, $\phi(0) = 0$ and ϕ'' is completely monotonous.

Moreover, Bernstein's lemma asserts that a fonction g continuous on $\mathbb{R}^+$, C^∞ on $]0, +\infty[$ is completely monotonous if and only if $\forall n \in \mathbb{N}, (-1)^n g^{(n)} \geq 0$. Injecting the definition of the exponents 2 in the evolution equation, one gets the following result.

Theorem 2. *Let $(\psi_t)_{t \geq 0}$ be a family of exponents of Levy processes of finite variance, such that $\forall \lambda \ t \mapsto \psi_t(\lambda)$ is differentiable and $\partial_t \psi_t$ is locally bounded. Then ψ_t characterizes an intrinsic statistical solution of our forced Burgers equation if and only if it satisfies the equation :*

$$\forall w \in \mathbb{R}, \partial_t \psi(t, w) = i \partial_w \psi(t, w) \psi(t, w) - \frac{w^2}{2}$$

If the family of exponents has negative jumps, the equation for ϕ is

$$(4) \quad \partial_t \phi(t, w) = -\partial_w \phi(t, w) \phi(t, w) + \frac{w^2}{2}$$

Proof. Suppose that u_t is a family of Levy processes of exponents $(t, w) \mapsto \psi(t, w)$. If v is a function in $\mathcal{D}_0$, the definition of ψ gives $\hat{\mu}_t(v) = \exp(\int_{\mathbb{R}} \psi(t, \int_x^{+\infty} v(y) dy) dx)$ (as can be seen by considering first simple functions). Hence, thanks to local boundedness, $\partial_t \hat{\mu}_t(v) = \hat{\mu}_t(v) \int_{\mathbb{R}} \partial_t \psi(t, \int_x^{+\infty} v(y) dy) dx = \hat{\mu}_t(v) \int_{\mathbb{R}} \partial_t \psi(t, w(x)) dx$ where we denote as before $w(x) = \int_x^{+\infty} v(y) dy$. The right hand side of (3) is more complicated to get. One needs to compute $\lim_{a \rightarrow -\infty, b \rightarrow +\infty} \int_E (\int_{\mathbb{R}} i \frac{1}{2} (u(x) - \frac{1}{b-a} \int_a^b u(y) dy)^2 v'(x) dx) \exp[i \int_{\mathbb{R}} u v] d\mu_t(u)$ in terms of ψ . This was done in [6] where it is shown that this limit is $i(\int_{\mathbb{R}} \partial_w \psi(t, w(x)) \psi(t, w(x)) dx) \hat{\mu}_t(v)$. The other term involves $\iint_{\mathbb{R} \times \mathbb{R}} v(x) v(y) A(x, y) dx dy$. We have to write it in terms of w : as noticed before, the specific form of A yields $\iint_{\mathbb{R} \times \mathbb{R}} v(x) v(y) A(x, y) dx dy = \int_{\mathbb{R}} w^2(x) dx$. Finally, we must have for all v in $\mathcal{D}_0$, $\int_{\mathbb{R}} \partial_t \psi(t, w(x)) dx = i \int_{\mathbb{R}} \partial_w \psi(t, w(x)) \psi(t, w(x)) dx - \frac{1}{2} \int_{\mathbb{R}} w^2(x) dx$. This is true if and only if $\forall w \in \mathbb{R}, \partial_t \psi(t, w) = i \partial_w \psi(t, w) \psi(t, w) - \frac{w^2}{2}$. $\square$

The stationary solution of equation (4) is $\phi(w) = \frac{w^{\frac{2}{3}}}{\sqrt{3}}$: this is the exponent of a Levy-stable process (which of course is not of finite variance: ϕ is not C^2 in 0). Speaking informally, this is an “invariant measure” of our forced Burgers equation.

Equation (4) can be solved using characteristics. Characteristics for (4) are curves $(\gamma_{x_0}(s), \lambda_{x_0}(s))$ that are solutions of the system

$$\begin{cases} \frac{d\gamma}{ds} &= \lambda(s) \\ \frac{d\lambda}{ds} &= \frac{\gamma(s)^2}{2} \end{cases}$$

with initial condition $\gamma_{x_0}(0) = x_0, \lambda_{x_0}(0) = \phi_0(x_0)$. Each x_0 corresponds to a different characteristics. To find the value of $\phi(x, t)$, one has to find x_0 , if it exists, such that $\gamma_{x_0}(t) = x$; if x_0 exists and is unique, then $\phi(x, t) = \lambda_{x_0}(t)$.

One can notice that system (3.1) describes the motion of a particle in a potential $V(x) = -\frac{x^3}{6}$, starting at x_0 with a velocity $\phi_0(x_0)$. Thanks to energy conservation, it is thus equivalent to $\lambda = \gamma', \gamma^3 - 3(\gamma')^2 = x_0^3 - 3\phi_0(x_0)^2, \gamma(0) = x_0$.

These remarks, and the preceding lemma characterizing Laplace exponents, are the basic ingredients of the next result.

Theorem 3. *Let ϕ_0 be the exponent of a Levy process of finite variance with negative jumps such that $\phi'_0(0) \geq 0$. Then the equation (4) with initial condition ϕ_0 admits a unique solution which is for all $t > 0$ the exponent of a homogeneous Levy process with finite variance and negative jumps. Such a Levy process is thus an intrinsic statistical solution of the randomly forced Burgers equation.*

If $\phi_0 = 0$, the solution is $\phi(x, t) = t^{-3} \sqrt{(xt^2)^3 - F(xt^2)^6}$ where F is the inverse of $x \mapsto 4^{\frac{1}{3}} x^2 \mathcal{P}(\omega_2 + \frac{x}{\sqrt{34^{\frac{1}{3}}}})$. Here $\mathcal{P}$ is the Weierstrass function and ω_2 its half-period.

We will start the proof by looking at the particular case $\phi_0 = 0$. It is instructive to see how things go, and it is anyway necessary to deal with it separately. The general case will be covered afterwards.

3.2. The particular case $\phi_0 = 0$. In this particular case (corresponding to u_0 constant), the system (3.1) is equivalent to $\lambda = \gamma', \gamma^3 - 3(\gamma')^2 = x_0^3, \gamma(0) = x_0$. The solution of this equation is $\gamma(s) = 4^{\frac{1}{3}}x_0\mathcal{P}(\omega_2 + \frac{\sqrt{x_0}}{\sqrt{34^{\frac{1}{3}}}}s)$ where $\mathcal{P}$ is the Weierstrass function solution of $y'^2 - 4y^3 + 1 = 0$ (with invariants $g_2 = 0$ and $g_3 = 1$), and $\omega_2 = \frac{\Gamma^3(\frac{1}{3})}{4\pi}$ is the half period of $\mathcal{P}$, satisfying $\mathcal{P}'(\omega_2) = 0$. γ is 1-1 from $[0, \frac{\sqrt{34^{\frac{1}{3}}}}{\sqrt{x_0}}\omega_2[$ to $[x_0, +\infty[$ (see [1]); it is C^∞ and strictly increasing on $]0, \frac{\sqrt{34^{\frac{1}{3}}}}{\sqrt{x_0}}\omega_2[$. In order to find $\phi(x, t)$, one needs to find x_0 such that $4^{\frac{1}{3}}x_0\mathcal{P}(\omega_2 + \frac{\sqrt{x_0}}{\sqrt{34^{\frac{1}{3}}}}t) = x$. Hence $x_0 = \frac{F(xt^2)^2}{t^2}$ where F is the inverse of $x \mapsto 4^{\frac{1}{3}}x^2\mathcal{P}(\omega_2 + \frac{x}{\sqrt{34^{\frac{1}{3}}}})$. F is continuous from $[0, +\infty[$ to $[0, \sqrt{44^{\frac{1}{3}}}\omega_2[$ and C^∞ on $]0, +\infty[$. Near 0, one has $4^{\frac{1}{3}}x^2\mathcal{P}(\omega_2 + \frac{x}{\sqrt{34^{\frac{1}{3}}}}) = x^2 + \frac{x^4}{4} + O(x^6)$, hence $F(x) = \sqrt{x} - \frac{x^{\frac{3}{2}}}{8} + o(x^2)$. Thus the solution of (4) with initial condition $\phi_0 = 0$ is $\phi(x, t) = t^{-3}\sqrt{(xt^2)^3 - F(xt^2)^6}$; the development of F near 0 ensures that $x \mapsto \phi(x, t)$ is C^2 on $[0, +\infty[$. The proof that ϕ'' is completely monotone will be done for the general case.

3.3. General case. As a preliminary remark, since ϕ_0'' is completely monotone, $\phi_0' \geq 0$ on $\mathbb{R}^+$, hence ϕ_0 is nonnegative and increasing on $\mathbb{R}^+$. Moreover, if there is an interval $I = [0, a[$ such that $\phi_0(w) = 0 \forall w \in I$, then $\phi_0'' = 0$ on I . But ϕ'' is decreasing (because $\phi_0^{(3)} \leq 0$) hence $\phi_0'' = 0$ on $\mathbb{R}^+$. Therefore $\phi_0 = 0$ on $\mathbb{R}^+$ or $\phi_0 > 0$ on $]0, +\infty[$. The particular case $\phi_0 = 0$ has been covered in the preceding section (except for complete monotonicity), hence we will assume here that $\phi_0 > 0$ on $\mathbb{R}^{+*}$.

Everywhere in this proof $\phi(x, t)', \phi(x, t)'', \phi(x, t)^{(n)}$ will refer to partial derivatives with respect to x .

As noticed before, the solution $\gamma_{x_0}(t)$ of the system (3.1) is the trajectory of a particle in a potential $V(x) = -\frac{x^3}{6}$, starting at x_0 with a velocity $\phi_0(x_0)$. Hence $\gamma_{x_0}(t)$ is a continuous increasing function of x_0 . More precisely, if $x_0 = 0$ $\gamma_0(t) = 0$, else if $x_0 \neq 0$,

$$\gamma_{x_0}(t) = x \Leftrightarrow t = \int_{x_0}^x \frac{dy}{\sqrt{y^3 - x_0^3 + 3\phi_0^2(x_0)}}$$

The function $h : \{(x, x_0)/0 < x_0 \leq x\} \rightarrow \mathbb{R}^+$, defined by $h(x, x_0) = \int_{x_0}^x \frac{dy}{\sqrt{y^3 - x_0^3 + 3\phi_0^2(x_0)}}$ is C^∞ ($\phi_0(x_0) > 0$, therefore this integral is well defined); $x_0 \mapsto h(x, x_0)$ is a diffeomorphism from $]0, x]$ into $\mathbb{R}^+$ hence for all $t > 0$ $\gamma_{x_0}(t) = x$ defines uniquely x_0 as a C^∞ function of (x, t) on $]0, +\infty[^2$, satisfying $\lim_{x \rightarrow 0} x_0(x, t) = 0$.

Hence the equation (4) admits as a solution for all t , $\phi(x, t) = \sqrt{x^3 - x_0(x, t)^3 + 3\phi_0^2(x_0)}$, $\phi(0, t) = 0$. $x \rightarrow \phi(x, t)$ is C^∞ on $]0, +\infty[$. We still have to check that ϕ'' is completely monotone and that ϕ is C^2 at 0.

Let us prove that ϕ'' is completely monotone. This proof will also work for $\phi_0 = 0$. The keypoint is to notice that $\frac{d}{dt}(\phi(\gamma_{x_0}(t), t)) = \lambda_{x_0}(t)\phi'(\gamma_{x_0}(t), t) + \frac{\partial \phi}{\partial t}(\gamma_{x_0}(t), t) = \phi\phi'(\gamma_{x_0}(t), t) + \frac{\partial \phi}{\partial t}(\gamma_{x_0}(t), t)$. Hence by deriving twice equation (4) with respect to w , and by setting $w = \gamma_{x_0}(t)$ one gets

$$\frac{d}{dt}(\phi''(\gamma_{x_0}(t), t)) + 3\phi'\phi''(\gamma_{x_0}(t), t) = 1$$

Thus, for every x_0 , $t \mapsto \phi''(\gamma_{x_0}(t), t)$ satisfies a differential inequality of the form $\frac{du}{dt} + 3fu \geq 0$ (where $f = \phi'(\gamma_{x_0}(t), t)$). Since $\phi_0'' \geq 0$, we deduce $\phi''(\gamma_{x_0}(t), t) \geq 0$ for all $t > 0$. Since it is true for all x_0 , $\phi''(x, t) \geq 0$ for all $x > 0, t > 0$.

One can now proceed by induction. Suppose $(-1)^n \phi^{(n)}(x, t) \geq 0$ for all $2 \leq n \leq N$. Then by deriving (4) $N + 1$ times one gets

$$\frac{d}{dt}(\phi^{(N+1)}(\gamma_{x_0}(t), t)) + (N+2)f\phi^{(N+1)}(\gamma_{x_0}(t), t) = -\frac{1}{2} \sum_{k=2}^N \binom{N+2}{k} \phi^k(\gamma_{x_0}(t), t) \phi^{(N+2-k)}(\gamma_{x_0}(t), t)$$

Now the induction hypothesis guarantees that the right hand side of this equation is of the sign of $(-1)^{N+1}$. Hence $(-1)^{N+1} \phi^{(N+1)}(\gamma_{x_0}(t), t)$ verifies $\frac{du}{dt} + (N+2)fu \geq 0$. One concludes as before that $(-1)^{N+1} \phi^{(N+1)}(x, t) \geq 0$.

Hence $\phi(x, t)$ is completely monotone.

Let us now prove that ϕ is C^2 at $x = 0$: we need to know how x_0 behaves near $x = 0$. In order to do this we will consider the solution of $y^3 - 3y'^2 + C(x_0) = 0, y(0) = x_0$, where $C(x_0) = 3\phi_0(x_0)^2 - x_0^3$. Its explicit expression depends on the sign of C . We know that $\lim_{x \rightarrow 0} x_0 = 0$. Hence there are really two cases that have to be dealt with : $\phi'_0(0) > 0$ and $\phi'_0(0) = 0$.

- If $\phi'_0(0) = 0$

ϕ_0 is C^2 at 0, hence $\phi_0(x_0) = k^2 x_0^2 + o(x_0^2)$ where k is a constant, and $C(x_0) = -x_0^3 + 3k^4 x_0^4 + o(x_0^4)$ is negative when x_0 is small enough. Hence $\gamma_{x_0}(t) = A\mathcal{P}(bt + c)$ where $A^3 = -4C(x_0)$, $b = \sqrt{\frac{A}{12}}$, $\mathcal{P}$ is as before the Weierstrass function with invariants $g_2 = 0, g_3 = 1$, and c verifies $A\mathcal{P}(c) = x_0$.

From the expansion $\mathcal{P}(\omega_2 + \epsilon) = 4^{-\frac{1}{3}} + 3.4^{-\frac{2}{3}}\epsilon^2 + o(\epsilon^2)$ we deduce

$$\begin{aligned} A &= 4^{\frac{1}{3}} x_0 (1 - k^4 x_0) + o(x_0^2) \\ b &= \sqrt{\frac{4^{\frac{1}{3}}}{12}} \sqrt{x_0} + o(\sqrt{x_0}) \\ c &= \omega_2 + \sqrt{\frac{4^{\frac{1}{3}}}{3}} k^2 \sqrt{x_0} + o(\sqrt{x_0}) \\ \gamma_{x_0}(t) &= x_0 + (k^2 t + \frac{t^2}{4}) x_0^2 + o(x_0^2) \\ x_0(x, t) &= x - (k^2 t + \frac{t^2}{4}) x^2 + o(x^2) \\ \phi(x, t) &= \sqrt{\frac{x^3 - x_0^3 + 3\phi_0(x_0)^2}{3}} = (k^2 + \frac{t}{2}) x^2 + o(x^2) \end{aligned}$$

Hence ϕ is C^2 at $x = 0$.

- If $\phi'_0(0) = a > 0$

Now $C(x_0) = 3a^2 x_0^2 + kx_0^3 + o(x_0^3)$, where k is a constant, is positive when x_0 is small enough. Hence $\gamma_{x_0}(t) = A\mathcal{Q}(bt + c)$ where $A^3 = 4C(x_0)$, $b = \sqrt{\frac{A}{12}}$, and $\mathcal{Q}$ is now the Weierstrass function with invariants $g_2 = 0, g_3 = -1$; and c must verify $A\mathcal{Q}(c) = x_0$. $\mathcal{Q}$ is related to $\mathcal{P}$ by $\mathcal{Q}(z) = -\mathcal{P}(iz)$ for all z in $\mathbb{C}$. $\mathcal{Q}$ admits a real zero z_1 , with $\mathcal{Q}'(z_1) = 1$ and $\mathcal{Q}''(z_1) = \mathcal{Q}'''(z_1) = 0$. It is not difficult to get the expansion $\mathcal{Q}(z_1 + \epsilon) = \epsilon + \frac{\epsilon^4}{2} + o(\epsilon^4)$, and :

$$\begin{aligned} A &= (12a^2)^{\frac{1}{3}} x_0^{\frac{2}{3}} (1 + \frac{k}{9a^2} x_0 + o(x_0)) \\ b &= (\frac{ax_0}{12})^{\frac{1}{3}} (1 + \frac{k}{18a^2} x_0 + o(x_0)) \\ c &= z_1 + (\frac{x_0}{12a^2})^{\frac{1}{3}} - a^{-\frac{8}{3}} (\frac{12^{-\frac{4}{3}}}{2} + k \frac{12^{-\frac{1}{3}}}{9}) x_0^{\frac{4}{3}} + o(x_0^{\frac{4}{3}}) \\ \gamma_{x_0}(t) &= x_0(1 + at) + (\frac{k}{18a} t + \frac{1}{24a^2} ((1 + at)^4 - 1)) x_0^2 + o(x_0^2) \\ x_0(x, t) &= \frac{x}{1 + at} - ((\frac{k}{18a} t + \frac{1}{24a^2} ((1 + at)^4 - 1)) \frac{x^2}{(1 + at)^3} + o(x^2)) \\ \phi(x, t) &= \sqrt{\frac{x^3 - x_0^3 + 3\phi_0(x_0)^2}{3}} = \frac{ax}{1 + at} \sqrt{1 + K_2 x + o(x)} \text{ where } K_2 = \frac{1 + 3(1 + at)^4 + 4k(1 + \frac{2at}{3})}{12a^2(1 + at)^2} \end{aligned}$$

Hence ϕ is here also C^2 at $x = 0$.

REFERENCES

- [1] M. ABRAMOVITZ and I.-A. STEGUN *Handbook of mathematical functions* Dover, 1965.
- [2] M. AVELLANEDA and W. E. Statistical properties of shocks in Burgers turbulence. *Commun. Math. Phys.*, 172(1):13-38, 1995.

- [3] J. BERTOIN. The inviscid Burgers equation with Brownian initial velocity. *Commun. Math. Phys.*, 193:397–406, 1998.
- [4] J.M. BURGERS. *The nonlinear diffusion equation*. Dordrecht: D. Reidel, 1974.
- [5] L. CARRARO and J. DUCHON. Solutions statistiques intrinsèques de l'équation de Burgers et processus de Lévy. *C. R. Acad. Sci., Paris, Sér. I*, 319:855–858, 1994.
- [6] L. CARRARO and J. DUCHON. Equation de Burgers avec conditions initiales à accroissements indépendants et homogènes. *Ann. Inst. Henri Poincaré Anal. Non Linéaire*, 15:431–458, 1998.
- [7] M.-L. CHABANOL and J. DUCHON. Markovian in space solutions of Burgers equation. *J. of Stat. Phys.*, 114:525–534, 2004.
- [8] W. E and E. vanden EIJNDEN. Statistical theory for the stochastic Burgers equation in the inviscid limit. *Comm. Pure Appl. Math.*, LIII:852–901, 2000.
- [9] L. FRACHEBOURG, P. A. MARTIN, and J. PIASECKI. Ballistic aggregation : a solvable model of irreversible many particle dynamics. *Physica A*, 279:69, 2000.
- [10] L. FRACHEBOURG and P.A. MARTIN. Exact statistical properties of the Burgers equation. *J. Fluid Mech.*, 417:323–349, 2000.
- [11] P. GROENEBOOM. Brownian motion with a parabolic drift and Airy functions. *Probab. Theory Related Fields*, 81:79–109, 1989.
- [12] D. REVUZ and M. YOR. *Continuous martingales and Brownian motion*. Springer, 1999.
- [13] R. ROBERT. Statistical hydrodynamics. In S. FRIEDLANDER and D. SERRE, editors, *Handbook of mathematical fluid mechanics*, volume 2. Elsevier, 2003.
- [14] Ya. G. SINAI. Statistics of shocks in solutions of the inviscid Burgers equation. *Commun. Math. Phys.*, 148:601–621, 1992.

INSTITUT DE MATHÉMATIQUES DE BORDEAUX, UMR 5251 CNRS-UNIVERSITÉ BORDEAUX1, FRANCE
E-mail address: Marie-Line.Chabanol@math.u-bordeaux1.fr

INSTITUT FOURIER (GRENOBLE), UMR 5582 CNRS-UNIVERSITÉ JOSEPH FOURIER, 100, RUE DES MATHÉMATIQUES,
 B.P. 74, 38041 SAINT-MARTIN D'HÈRES CEDEX, FRANCE
E-mail address: Jean.Duchon@ujf-grenoble.fr